TMA GENERIC QUALITY SCREENS

1. Adequacy of Discharge Planning

No documentation of discharge planning or appropriate follow-up care with consideration of physical, emotional and mental status needs at time of discharge.

2. Medical Stability of the Patient

a. Blood Pressure:

Adults (within 24 hours of discharge): systolic less than 85 or greater than 180; diastolic less than 50 or greater than 110 (obstetrical patients a diastolic of 100)

.

<u>Children</u> (prior to discharge, EXCEPT newborns):

<u>Age</u>	Systolic Range	<u>Diastolic Range</u>
0 - 6	78-114	46-80
6 - 12	82-132	50-88
12 - 18	98-146	60-95

b. <u>Temperature within 24 hours of discharge greater than:</u>

- 101 degrees Fahrenheit (38.3 Centigrade) oral, or
- 102 degrees Fahrenheit (38.9 Centigrade) rectal, or
- 100 degrees Fahrenheit (37.8 Centigrade) axillary

c. <u>Pulse within 24 hours of discharge:</u>

Adults: less than 50 (or 45 if patient is on a Beta blocker or an athlete) or greater than 120.

Attachment J-2 Page 1

Children:

Age	Below	Above
0 - 3 months	60	200
3 months - 2 years	70	160
2 - 6 years	75	120
6 - 18 years	65	110

- d. Abnormal diagnostic findings which are not addressed and resolved or, where the record does not explain why they are not resolved.
- e. Purulent or bloody drainage of wound or open area within 24 hours prior to discharge (excludes from review normal discharges associated with post-partum recovery).
 - f. Pediatric respiratory rate within 24 hours of discharge:

<u>Age</u>	Rate above
0 - 2	60/minute
2 - 18	32/minute

or physician documented respiratory distress.

g. Weight less than 2.0kg for pediatric newborn or premature infants.

3. Deaths

- a. During or following any surgery performed during the current admission.
- b. Following return to intensive care, coronary care or other special unit within 24 hours of being transferred out.
 - c. Other unexpected death.

4. Nosocomial Infection

Attachment J-2 Page 2

(Hospital Acquired Infection)

5. Unscheduled Return to Surgery

Within same admission for same condition as previous surgery or to correct operative problem.

6. Trauma Suffered in the Hospital

- a. Unplanned surgery which includes, but is not limited to removal or repair of a normal organ or body part (i.e., surgery not addressed specifically in the operative consent).
 - b. Fall.
 - c. Serious complications of anesthesia.
 - d. Any transfusion error or serious transfusion reaction.
- e. Hospital acquired decubitus ulcer and/or deterioration of an existing decubitus.
- f. Medication error or adverse drug reaction (1) with serious potential for harm or (2) resulting in measures to correct.
- g. Care or lack of care resulting in serious or potentially serious complications.

Attachment J-2 Page 3